



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{\Delta_m}{\Delta_{m+1}} = \sqrt{\left(1 - \frac{1}{m^2}\right)} \dots (4).$$

II. Solution by G. B. M. ZERR, A. M., Ph. D., Vice President and Professor of Mathematics in the Texarkana College, Texarkana, Arkansas.

Let p be the excess of pressure, on the assumption that the external pressure is constant.

Let u =original volume, then when the sphere is n times its original volume, the work done in this electrification must be

$$p(n-1)u, \text{ (see Minchin's Statics).}$$

Let v be the potential, Q the charge of this electrification, r =radius of sphere, σ =electrical density.

$$\therefore \text{energy of electrification} = \frac{1}{2}vQ. \text{ But } v = \frac{Q}{r} \text{ and } Q = 4\pi r^2 \sigma.$$

$$\therefore p(n-1)u = \frac{1}{2} \frac{Q^2}{r} = 8\pi^2 r^3 \sigma^2. \text{ But } \frac{4}{3}\pi r^3 = nu.$$

$$\therefore p(n-1)u = 6n\pi\sigma^2 u.$$

$$\therefore p(n-1) = 6n\pi\sigma^2.$$

Similarly, when the sphere becomes $(m+1)$ times its original size we get, if σ_1 is the density, $pm = 6(m+1)\pi\sigma_1^2$.

$$\therefore \frac{\sigma}{\sigma_1} = \sqrt{\frac{(m+1)(n-1)}{mn}} = \sqrt{\frac{n^2-1}{m^2}} \text{ if } n=m.$$

27. Proposed by G. B. M. ZERR, A. M., Ph. D., Vice President and Professor of Mathematics in the Texarkana College, Texarkana, Arkansas.

One thousand balls, each having a mass of 10 grams, and each moving with a velocity of 10 kilometers per second, are confined in a certain space with elastic walls. Into the same space are now introduced one thousand balls each of 1000 grams mass, and moving with a velocity each of 10 kilometers per second; collisions take place, and finally, after a number of encounters, the average kinetic energy of each of the two thousand balls is the same. Show that this is 5.75(10)¹¹ in the centimeter-gram-system.

Solution by O. W. ANTHONY, Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland and the PROPOSER.

By Avogadro's hypothesis and kinetic theory, if M, M_1 are the masses and V, V_1 the velocities of two balls, then $\frac{1}{2}MV^2$ and $\frac{1}{2}M_1V_1^2$ are their respective kinetic energies. As the average kinetic energy of each ball is equal, we get $\frac{1}{2}MV^2 = \frac{1}{2}M_1V_1^2$.

$$\therefore \text{The average kinetic energy} = \frac{1}{2}(\frac{1}{2}MV^2 + \frac{1}{2}M_1V_1^2) = E.$$

$$\therefore E = \frac{1}{4}(MV^2 + M_1V_1^2).$$

$$\text{Let } V = V_1, \text{ then } E = \frac{1}{4}(M + M_1)V^2.$$

$$V = 10 \text{ kilometers} = 10^6 \text{ centimeters.}$$

$$M = 10 \text{ grams, } M_1 = 100 \text{ grams.}$$

$$\therefore E = \frac{1}{4}(10 + 100)10^{12} = \frac{1}{4}11 \times 10^{13} = 2.75 \times 10^{13} = 2.75(10)^{13}.$$

Also solved by F. P. MATZ.